

# OTHER SYMPTOMS TO LOOK OUT FOR

Below are other symptoms that you can have when you take Opdivo®. We are very concerned about symptoms we have marked with a check mark. Please call the office if you develop any of these symptoms, and consider going directly to the Emergency Room if you have any of the red-flag symptoms.

| ✓ | Symptoms  | Red flags (Signs that something serious is going on that needs immediate attention)  |
|---|---|--|
|   | Appetite changes (not feeling hungry or wanting to eat) | If you have no appetite plus you feel faint, pass out, and/or have trouble thinking (cloudy thinking, confusion)   |
|   | Chest pain  | Any chest pain should be evaluated immediately   |
|   | Fatigue (feeling weak or tired)                         | If you faint or also have shortness of breath or chest pain  |
|   | Fever   | <ul style="list-style-type: none"> <li>• If you have a fever over 102.5°F</li> <li>• If you also have breathing problems</li> </ul>  |
|   | Headaches   | <ul style="list-style-type: none"> <li>• If you have a very painful headache (the “worst headache of your life”), head pain that doesn’t go away after a few hours, or head pain with fever, neck pain or stiffness, and/or rash</li> <li>• If you have head pain and changes in vision</li> </ul> |
|   | Nausea  | If you have uncontrolled nausea and it is limiting your ability to eat or drink  |
|   | Shortness of breath (trouble breathing)                 | <ul style="list-style-type: none"> <li>• If you have any trouble breathing or worsening of your existing symptoms</li> <li>• If you have shortness of breath plus chest pain and/or fainting</li> <li>• If you have shortness of breath and swelling of the legs or leg pain</li> </ul>            |